Web Images Videos Maps News Shopping Gmail more⊸

Sign in



protein dot-blot TCA methanol SDS ×

arch instant is on ▼

About 5,710 results (0.10 seconds)

Advanced search

Immun-Blot® PVDF Membrane for Protein

Blotting Instruction Manual File Format: PDF/Adobe Acrobat

Dot blotting requires special precautions to insure that **protein** is bound to the **PVDF** mem-... Blot **PVDF** membrane is not affected by the addition of this amount of **SDS**. ... **Methanol** should be analytical reagent grade, as metallic ...

www.big-

rad.com/webroot/web/pdf/\sr/literature/Bulletin 4006127A.pdf

Western Blotting PVDF Membranes

Western **blot** transfer from an **SDS**-PAGE gel is a method that can be used to detect individual ... **Protein dot** or slot **blots**, N terminal **protein** sequencing Destain for 3 minutes with Coomassie Blue Destain I (80% **methanol**; 10% acetic acid). Western **blotted proteins** on **PVDF** membranes can be detected by ...

www.pall.com/laboratory_48923.asp - Cached - Similar

Immun-Blot® PVDF Membrane for Protein Blotting Instruction Manual

Dot blotting requires special precautions to insure that **protein** is bound to ... This stabilizes the effect of **methanol** in stripping **SDS** off the **proteins**. ...

www.scribd.com/.../Immun-Blot®-PVDF-Membrane-for-Protein-Blotting-Instruction-Manual - Cached - Similar

Purifying proteins for proteomics: a laboratory manual - Google Books Result

Richard J. Simpson - 2004 - Medicai - 801 pages
Polyvinyldifluoride (PVDF) membranes are hydrophobic and
offer a uniformly ... Reference protein solution (1 mg/ml)
Semiquantitative dot blotting requires ...
books.google.com/books?isbn=0879696966...

Quantitative filtration-blotting of protein in the presence of ...

by T Nishibu - 2003 - Cited by 8 - Related articles

A variety of **proteins** were blotted using a **dot-blot** apparatus with a **PVDF** ... up by fitting a sheet of hydrated **PVDF** membrane with a 0.1-µm pore size, ... to melt the membrane, and ethanol could be replaced with **methanol**. ... varying **SDS** concentrations in the blotting solution containing 2.5% **TCA** and 45% ethanol. ...

linkinghub.elsevier.com/retrieve/pii/S0003269703002550

Everything

Images

Videos

News

Shopping

Books

More

Alexandria, VA

Change location

Show search tools

Estimating Protein Concentration by Dot Blotting of Multiple ...

The **dot blot** involves direct application of dilutions of test **protein** to a ... Because **TCA** is extremely hygroscopic and thus difficult to weigh ... However, **proteins** bind poorly to nitrocellulose in the presence of detergents such as **SDS** If using a **PVDF** membrane, wet the membrane in 100% **methanol** for 15 ...

cshprotocols.cshlp.org/cgi/content/full/2006/1/pdb.prot4253?...
- Similar

[PDF]

T- cell leukemias and lymphomas Characterization of a new T ...

File Format: PDF/Adobe Acrobat - Quick View by WF Cassano - 1990 - Related articles ... Blue in 45% **methanol**, 10% acetic acid in water and destained in 25% **methanol**, 7.5% acetic acid in water. ... was then electroblotted to **PVDF** membranes (Millipore, Bedford, ... **SDS**, was applied under suction to the 96-well **dot blot** apparatus ... Sepharose beads (Vector) before

bloodjournal.hematologylibrary.org/cgi/reprint/75/7/1557.pdf

Western blotting Methods, Protocols and Troubleshootings

performing the TCA-1 protein dot ...

SYPRO RUBY **protein blot** staining and Subsequent Western **blot** - (reply: 3) ... western transfer efficiency.. with or without **methanol** / **SDS?** ... Nitrocellulose or **PVDF** membrane for western **blot** - which one? Problem with **TCA** and western **blot** - (reply: 3); Concentrating **protein** in cerebrospinal fluid for Western ...

www.protocol-online.org/biology.../Western-biotting.html - Cached - Similar

Molecular Cloning, Sequence Analysis, and Immune Recognition of ...

by S dittapalapong - 2008 - Cited by 3 - Related articles formed by induction with 0.5% **methanol** every ... Coomassie blue or transferred onto **PVDF** at. 400 mAmp for 5 h at 4 ... **dot blot** analysis of recombinant **proteins** secreted into supernatant at day 2 of wild type (wt) ... 10% **SDS**-PAGE analysis of crude **protein** harvested at days 0, 1, 2, 3, and 4 (lanes 1–5, ...

onlinelibrary.wiley.com/doi/10.1196/annals.1428.059/pdf

ppp Matchmaker™ Monoclonal Antibodies

File Format: PDF/Adobe Acrobat - Quick View May 1, 2009 ... performed on a **dot blot** if an appropriate, purified control **protein** Nitrocellulose (0.2-µm pore size, Schliecher & Schuell; or **PVDF** ... 200 ml of 100% **methanol** deionized H2O ... Blot **proteins** onto the nitrocellulose paper according to the **SDS** method (Panel A) or the **TCA** method (Panel B). ...

www.clontech.com/images/pt/PT1029-1.pdf - Similar

1 2 3 4 5 6 7 8 9 10 Next

Search Help Give us feedback

Google Home - Advertising Programs - Business Solutions - Privacy - About Google